

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
7 November 2002 (07.11.2002)

PCT

(10) International Publication Number  
WO 02/088178 A2

(51) International Patent Classification<sup>7</sup>: C07K 14/315

(21) International Application Number: PCT/CA02/00664

(22) International Filing Date: 2 May 2002 (02.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/287,712 2 May 2001 (02.05.2001) US

(71) Applicant (for all designated States except US): SHIRE  
BIOCHEM INC. [CA/CA]; 275 Armand-Frappier Boulevard,  
Laval, Québec H7V 4A7 (CA).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MARTIN, Denis  
[CA/CA]; 4728-G rue Gaboury, St-Augustin-de-Des-  
maures, Québec G3A 1E9 (CA). HAMEL, Josée  
[CA/CA]; 2401 Maritain, Sillery, Québec G1T 1N6 (CA).  
BRODEUR, Bernard, R. [CA/CA]; 2401 rue, Sillery,  
Québec G1T 1N6 (CA). RIOUX, Stéphane [CA/CA]; 869  
Avenue Des Pinsons, Beauport, Québec G1E 1J3 (CA).  
BOYER, Martine [CA/CA]; 3924 Rue Louise-Fiset,  
Ste-Foy, Québec G1X 4T7 (CA).

(74) Agents: MORROW, Joy, D. et al.; Smart & Biggar, P.O.  
Box 2999, Station D, 900-55 Metcalfe Street, Ottawa, On-  
tario K1P 5Y6 (CA).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).

Published:

— without international search report and to be republished  
upon receipt of that report

[Continued on next page]

(54) Title: ANTIGENS OF GROUP B STREPTOCOCCUS AND CORRESPONDING DNA FRAGMENTS

```
1 MTKKHLKTLA LALTTVSVVT YSQEVYGLER EBSVKQEOQT SASEDDWPEE
DNERKTNVSK
61 ENSTVDETVS DLPDGNNSNN SSKTESVVS DPKQVPKAKP EVTQEASNSS
NDASKVEVPR
121 QDTASKKBTL ETSTWEAKDF VTRGDTLVGF SKSGINKLSQ TSHLVLP SHA
ADGTQLTOVA
181 SFAFTPDKKT AIAEYTSRLG ENGKPSRLDI DQKEIIDEGE IFNAYQLTKL
TIPNGYKSIG
241 QDAFVDNKN IAEVNLPESE TISDYAFAMH SLKQVKLPDN LKVGELAF
DNQIGGKLYL
301 PRHLIKLAER AFKSNRIQTV EPLGSKLVKI GEASFQDNNL RNVMLPDGLE
KIESEAF TGN
361 PGDEHYNNQV VLRTRTQGNP HQLATENTYV NPDKSLWRAT PDMYTKWLE
EDFTYQKNSV
421 TGFSNKGK LQK VRRNKNLEIP KQNGITITE IGDNAFRNVD FQSKTLRKYD
LEEIKLPSTI
481 RKIGAFAPQS NNKLSFEASE DLEBIKEGAF MNNRIGTLDL KDKLIKIGDA
APHINHIYAI
541 VLPESVQBIG RSAFRQNGAL HLMPIGNKVK TIGEMAFLSN KLESVNLSEQ
KQLKTLEVQA
601 FSDNALSEVV LPPNLQTIRE EAFKRNHLKE VKGSSTLSQI TFNAPQNDG
DKRFGKVVV
661 RTHNNSHMLA DGERFIIDPD KLSSTMDLE KVLKIIEGLD YSTLRQTTQT
QPREMTTAGK
721 ALLSKSNLRQ GEKQKFLQEA QFFLGRVDLD KAIKAKAKAL VTKKATKNHG
LLERSINKAV
781 LAYNNSAIKK ANVKRLEKEL DLLTDLVRGK GPLAQATMVQ GVVYLLKTPLP
LPEYYIGLNV
841 YFDKSGKLIY ALDMSDTIGE GQKDAYGNPI LNVEDENEGY HTLAVATLAD
YEGLYIKDIL
901 NSSLDKIKAI RQIPLAKYHR LGIFQAIRNA AAEADRLLPK TPKGYLNEVP
NVRKKQMEKN
961 LKPVYDYKTP I FNKALPNEKV DGDRAAKGHN INAETNNSVA VTPIRSEQQL
HKSQSDVNL
1021 QTSSKNNFIY EILGYVSLCL LPLVTAGKKG KRARK*
//
```

(57) Abstract: The present invention relates to antigens, more particularly antigens of Group B Streptococcus (GBS) (*S. agalactiae*) which may be useful to prevent, diagnose and/or treat streptococcal infections.

WO 02/088178 A2



---

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

ANTIGENS OF GROUP B STREPTOCOCCUS and CORRESPONDING DNA  
FRAGMENTS

5 FIELD OF THE INVENTION

The present invention is related to antigens, more particularly polypeptides of Group B Streptococcus (GBS) (S. agalactiae) which may be used to prevent, diagnose, and/or treat GBS infections.

10

BACKGROUND OF THE INVENTION

Streptococcus are gram (+) bacteria that are differentiated by group specific carbohydrate antigens A through O found on their cell surface. Streptococcus groups are further  
15 distinguished by type-specific capsular polysaccharide antigens. Several serotypes have been identified for the GBS: Ia, Ib, II, III, IV, V, VI, VII and VIII. GBS also contains antigenic proteins known as "C-proteins" (alpha, beta, gamma and delta), some of which have been cloned.

20

Although GBS is a common component of the normal human vaginal and colonic flora this pathogen has long been recognized as a major cause of infections in neonates or infants, expectant mothers, some non-pregnant adults as well  
25 as mastitis in dairy herds. Expectant mothers exposed to GBS are at risk of postpartum infection and may transfer the infection to their baby as the child passes through the birth canal.

30 GBS infections in infants are restricted to very early infancy. Approximately 80% of infant infections occur in the first days of life, so-called early-onset disease. Late-onset infections occur in infants between 1 week and 2 to 3 months of age. Clinical syndromes of GBS disease in  
35 newborns include sepsis, meningitis, pneumonia, cellulitis, osteomyelitis, septic arthritis, endocarditis and epiglottitis. In addition to acute illness due to GBS, which

is itself costly, GBS infections in newborns can result in death, disability, and, in rare instances, recurrence of infection. Although the organism is sensitive to antibiotics, the high attack rate and rapid onset of sepsis in neonates and meningitis in infants results in high morbidity and mortality.

Among pregnant women, GBS causes clinical illness ranging from mild urinary tract infection to life-threatening sepsis and meningitis, including also osteomyelitis, endocarditis, amniotitis, endometritis, wound infections (postcesarean and postepisiotomy), cellulitis, fasciitis.

Among non-pregnant adults, the clinical presentations of invasive GBS disease most often take the form of primary bacteremia but also skin or soft tissue infection, pneumonia, urosepsis, endocarditis, peritonitis, meningitis, empyema. Skin or soft tissue infections include cellulitis, infected peripheral ulcers, osteomyelitis, septic arthritis and decubiti or wound infections. Among people at risk, there are debilitated hosts such as people with a chronic disease such as diabetes mellitus and cancer, or elderly people.

GBS infections can also occur in animals and cause mastitis in dairy herds.

Type-specific polysaccharides have proven to be poorly immunogenic in hosts and are restricted to the particular serotype from which the polysaccharide originates. Further, capsular polysaccharide elicit a T cell independent response i.e. no IgG production. Consequently capsular polysaccharide antigens are unsuitable as a vaccine component for protection against GBS infection.

Others have focused on the C-protein beta antigen which demonstrated immunogenic properties in mice and rabbit models. This protein was found to be unsuitable as a human vaccine because of its undesirable property of interacting

with high affinity and in a non-immunogenic manner with the Fc region of human IgA. The C-protein alpha antigen is rare in type III serotypes of GBS which is the serotype responsible for most GBS mediated conditions and is therefore of little use as a vaccine component.

There remains an unmet need for GBS antigens that may be used as vaccine components for the prophylaxis and/or therapy of GBS infection.

#### SUMMARY OF THE INVENTION

According to one aspect, the present invention provides an isolated polynucleotide encoding a polypeptide having at least 80% identity to a second polypeptide comprising a sequence chosen from SEQ ID Nos : 2, or fragments or analogs thereof.

According to one aspect, the present invention relates to polypeptides comprising SEQ ID No : 2 or fragments or analogs thereof.

In other aspects, there are provided polypeptides encoded by polynucleotides of the invention, pharmaceutical composition, vectors comprising polynucleotides of the invention operably linked to an expression control region, as well as host cells transfected with said vectors and processes for producing polypeptides comprising culturing said host cells under conditions suitable for expression.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 represents the DNA sequence of BVH-A4 gene from serotype III Group B streptococcus strain COH1; (SEQ ID NO: 1). The underlined portion of the sequence represents the region coding for the leader peptide.

Figure 2 represents the amino acid sequence of BVH-A4 protein from serotype III Group B streptococcus strain COH1;

(SEQ ID NO: 2). The underlined sequence represents the 22 amino acid residues leader peptide.

5 DETAILED DESCRIPTION OF THE INVENTION

The present invention provides purified and isolated polynucleotides, which encode Streptococcal polypeptides that may be used to prevent, diagnose and/or treat Streptococcal infection.

10

According to one aspect, the present invention provides an isolated polynucleotide encoding a polypeptide having at least 80% identity to a second polypeptide comprising a sequence chosen from SEQ ID NO: 2 or fragments or analogs thereof.

15

According to one aspect, the present invention provides an isolated polynucleotide encoding a polypeptide having at least 90% identity to a second polypeptide comprising a sequence chosen from SEQ ID NO: 2 or fragments or analogs thereof.

20

According to one aspect, the present invention provides an isolated polynucleotide encoding a polypeptide having at least 95% identity to a second polypeptide comprising a sequence chosen from SEQ ID NO: 2 or fragments or analogs thereof.

25

According to one aspect, the present invention relates to polypeptides comprising SEQ ID No : 2 or fragments or analogs thereof.

30

According to one aspect, the present invention provides a polynucleotide encoding an epitope bearing portion of a

polypeptide comprising SEQ ID NO: 2 or fragments or analogs thereof.

According to one aspect, the present invention relates to  
5 epitope bearing portions of a polypeptide comprising SEQ ID NO: 2 or fragments or analogs thereof.

According to one aspect, the present invention provides an isolated polynucleotide encoding a polypeptide having at  
10 least 80% identity to a second polypeptide comprising SEQ ID NO: 2.

According to one aspect, the present invention provides an isolated polynucleotide encoding a polypeptide having at  
15 least 90% identity to a second polypeptide comprising SEQ ID NO: 2.

According to one aspect, the present invention provides an isolated polynucleotide encoding a polypeptide having at  
20 least 95% identity to a second polypeptide comprising SEQ ID NO: 2.

According to one aspect, the present invention relates to polypeptides comprising SEQ ID No : 2.  
25 According to one aspect, the present invention provides a polynucleotide encoding an epitope bearing portion of a polypeptide comprising SEQ ID NO: 2.

According to one aspect, the present invention relates to  
30 epitope bearing portions of a polypeptide comprising SEQ ID NO: 2.

According to one aspect, the present invention provides an isolated polynucleotide comprising a polynucleotide chosen from:

- 5 (a) a polynucleotide encoding a polypeptide having at least 80% identity to a second polypeptide comprising a sequence chosen from: SEQ ID NO: 2 or fragments or analogs thereof;
- (b) a polynucleotide encoding a polypeptide having at least 95% identity to a second polypeptide comprising a  
10 sequence chosen from: SEQ ID NO: 2 or fragments or analogs thereof;
- (c) a polynucleotide encoding a polypeptide comprising a sequence chosen from: SEQ ID NO: 2 or fragments or analogs thereof;
- 15 (d) a polynucleotide encoding a polypeptide capable of generating antibodies having binding specificity for a polypeptide comprising a sequence chosen from: SEQ ID NO: 2 or fragments or analogs thereof;
- (e) a polynucleotide encoding an epitope bearing portion of a polypeptide comprising a sequence chosen from SEQ ID  
20 NO: 2 or fragments or analogs thereof;
- (f) a polynucleotide comprising a sequence chosen from SEQ ID NO: 1 or fragments or analogs thereof;
- (g) a polynucleotide that is complementary to a  
25 polynucleotide in (a), (b), (c), (d), (e) or (f).

According to one aspect, the present invention provides an isolated polynucleotide comprising a polynucleotide chosen from:

- 30 (a) a polynucleotide encoding a polypeptide having at least 80% identity to a second polypeptide comprising a sequence chosen from: SEQ ID NO: 2;



- (b) a polynucleotide encoding a polypeptide having at least 95% identity to a second polypeptide comprising a sequence chosen from: SEQ ID NO: 2;
- (c) a polynucleotide encoding a polypeptide comprising a sequence chosen from: SEQ ID NO: 2;
- (d) a polynucleotide encoding a polypeptide capable of raising antibodies having binding specificity for a polypeptide comprising a sequence chosen from: SEQ ID NO: 2;
- (e) a polynucleotide encoding an epitope bearing portion of a polypeptide comprising a sequence chosen from SEQ ID NO: 2;
- (f) a polynucleotide comprising a sequence chosen from SEQ ID NO: 1;
- (g) a polynucleotide that is complementary to a polynucleotide in (a), (b), (c), (d), (e) or (f).

According to one aspect, the present invention provides an isolated polypeptide comprising a polypeptide chosen from:

- (a) a polypeptide having at least 80% identity to a second polypeptide comprising SEQ ID NO: 2, or fragments or analogs thereof;
- (b) a polypeptide having at least 95% identity to a second polypeptide comprising SEQ ID NO: 2, or fragments or analogs thereof;
- (c) a polypeptide comprising SEQ ID NO: 2, or fragments or analogs thereof;
- (d) a polypeptide capable of raising antibodies having binding specificity for a polypeptide comprising SEQ ID NO: 2, or fragments or analogs thereof;
- (e) an epitope bearing portion of a polypeptide comprising SEQ ID NO: 2, or fragments or analogs thereof;
- (f) the polypeptide of (a), (b), (c), (d), (e) or (f) wherein the N-terminal Met residue is deleted;

- (g) the polypeptide of (a), (b), (c), (d), (e) or (f)  
wherein the secretory amino acid sequence is deleted.

According to one aspect, the present invention provides an  
5 isolated polypeptide comprising a polypeptide chosen from:

- (a) a polypeptide having at least 80% identity to a second polypeptide comprising SEQ ID NO: 2;
- (b) a polypeptide having at least 95% identity to a second polypeptide comprising SEQ ID NO: 2;
- 10 (c) a polypeptide comprising SEQ ID NO: 2;
- (d) a polypeptide capable of raising antibodies having binding specificity for a polypeptide comprising SEQ ID NO: 2;
- (e) an epitope bearing portion of a polypeptide comprising  
15 SEQ ID NO: 2;
- (f) the polypeptide of (a), (b), (c), (d), (e) or (f) wherein the N-terminal Met residue is deleted;
- (g) the polypeptide of (a), (b), (c), (d), (e) or (f) wherein the secretory amino acid sequence is deleted.

20

Those skilled in the art will appreciate that the invention includes DNA molecules, i.e. polynucleotides and their complementary sequences that encode analogs such as mutants, variants, homologues and derivatives of such polypeptides,  
25 as described herein in the present patent application. The invention also includes RNA molecules corresponding to the DNA molecules of the invention. In addition to the DNA and RNA molecules, the invention includes the corresponding polypeptides and monospecific antibodies that specifically  
30 bind to such polypeptides.

In a further embodiment, the polypeptides in accordance with the present invention are antigenic.

In a further embodiment, the polypeptides in accordance with the present invention are immunogenic.

- 5 In a further embodiment, the polypeptides in accordance with the present invention can elicit an immune response in a host.

10 In a further embodiment, the present invention also relates to polypeptides which are able to raise antibodies having binding specificity to the polypeptides of the present invention as defined above.

15 An antibody that "has binding specificity" is an antibody that recognizes and binds the selected polypeptide but which does not substantially recognize and bind other molecules in a sample, e.g., a biological sample, which naturally includes the selected peptide. Specific binding can be measured using an ELISA assay in which the selected  
20 polypeptide is used as an antigen.

In accordance with the present invention, "protection" in the biological studies is defined by a significant increase in the survival curve, rate or period. Statistical analysis  
25 using the Log rank test to compare survival curves, and Fisher exact test to compare survival rates and numbers of days to death, respectively, might be useful to calculate P values and determine whether the difference between the two groups is statistically significant. P values of 0.05 are  
30 regarded as not significant.

In an additional aspect of the invention there are provided antigenic/immunogenic fragments of the polypeptides of the invention, or of analogs thereof.

5 The fragments of the present invention should include one or more such epitopic regions or be sufficiently similar to such regions to retain their antigenic/immunogenic properties. Thus, for fragments according to the present invention the degree of identity is perhaps irrelevant,  
10 since they may be 100% identical to a particular part of a polypeptide or analog thereof as described herein. The present invention further provides fragments having at least 10 contiguous amino acid residues from the polypeptide sequences of the present invention. In one embodiment, at  
15 least 15 contiguous amino acid residues. In one embodiment, at least 20 contiguous amino acid residues.

The skilled person will appreciate that analogs of the polypeptides of the invention will also find use in the  
20 context of the present invention, i.e. as antigenic/immunogenic material. Thus, for instance proteins or polypeptides which include one or more additions, deletions, substitutions or the like are encompassed by the present invention.

25 As used herein, "fragments", "analogs" or "derivatives" of the polypeptides of the invention include those polypeptides in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid  
30 residue (preferably conserved) and which may be natural or unnatural. In one embodiment, derivatives and analogs of polypeptides of the invention will have about 80% identity with those sequences illustrated in the figures or fragments

thereof. That is, 80% of the residues are the same. In a further embodiment, polypeptides will have greater than 80% identity. In a further embodiment, polypeptides will have greater than 85% identity. In a further embodiment, polypeptides will have greater than 90% identity. In a further embodiment, polypeptides will have greater than 95% identity. In a further embodiment, polypeptides will have greater than 99% identity. In a further embodiment, analogs of polypeptides of the invention will have fewer than about 20 amino acid residue substitutions, modifications or deletions and more preferably less than 10.

These substitutions are those having a minimal influence on the secondary structure and hydropathic nature of the polypeptide. Preferred substitutions are those known in the art as conserved, i.e. the substituted residues share physical or chemical properties such as hydrophobicity, size, charge or functional groups. These include substitutions such as those described by Dayhoff, M. in Atlas of Protein Sequence and Structure 5, 1978 and by Argos, P. in EMBO J. 8, 779-785, 1989. For example, amino acids, either natural or unnatural, belonging to one of the following groups represent conservative changes:

ala, pro, gly, gln, asn, ser, thr, val;

cys, ser, tyr, thr;

val, ile, leu, met, ala, phe;

lys, arg, orn, his;

and phe, tyr, trp, his.

The preferred substitutions also include substitutions of D-enantiomers for the corresponding L-amino acids.

In an alternative approach, the analogs could be fusion proteins, incorporating moieties which render purification

easier, for example by effectively tagging the desired polypeptide. It may be necessary to remove the "tag" or it may be the case that the fusion polypeptide itself retains sufficient antigenicity to be useful.

5

The percentage of homology is defined as the sum of the percentage of identity plus the percentage of similarity or conservation of amino acid type.

10 In one embodiment, analogs of polypeptides of the invention will have about 80% identity with those sequences illustrated in the figures or fragments thereof. That is, 80% of the residues are the same. In a further embodiment, polypeptides will have greater than 85% identity. In a  
15 further embodiment, polypeptides will have greater than 90% identity. In a further embodiment, polypeptides will have greater than 95% identity. In a further embodiment, polypeptides will have greater than 99% identity. In a further embodiment, analogs of polypeptides of the invention  
20 will have fewer than about 20 amino acid residue substitutions, modifications or deletions and more preferably less than 10.

In one embodiment, analogs of polypeptides of the invention  
25 will have about 80% homology with those sequences illustrated in the figures or fragments thereof. In a further embodiment, polypeptides will have greater than 85% homology. In a further embodiment, polypeptides will have greater than 90% homology. In a further embodiment,  
30 polypeptides will have greater than 95% homology. In a further embodiment, polypeptides will have greater than 99% homology. In a further embodiment, analogs of polypeptides

of the invention will have fewer than about 20 amino acid residue substitutions, modifications or deletions and more preferably less than 10.

5 One can use a program such as the CLUSTAL™ program to compare amino acid sequences. This program compares amino acid sequences and finds the optimal alignment by inserting spaces in either sequence as appropriate. It is possible to calculate amino acid identity or homology for an optimal  
10 alignment. A program like BLASTx™ will align the longest stretch of similar sequences and assign a value to the fit. It is thus possible to obtain a comparison where several regions of similarity are found, each having a different score. Both types of identity analysis are contemplated in  
15 the present invention.

In an alternative approach, the analogs or derivatives could be fusion polypeptides, incorporating moieties which render purification easier, for example by effectively tagging the  
20 desired protein or polypeptide, it may be necessary to remove the "tag" or it may be the case that the fusion polypeptide itself retains sufficient antigenicity to be useful.

25 It is well known that is possible to screen an antigenic polypeptide to identify epitopic regions, i.e. those regions which are responsible for the polypeptide's antigenicity or immunogenicity. Methods for carrying out such screening are well known in the art. Thus, the fragments of the present  
30 invention should include one or more such epitopic regions or be sufficiently similar to such regions to retain their antigenic/immunogenic properties.

Thus, for fragments according to the present invention the degree of identity is perhaps irrelevant, since they may be 100% identical to a particular part of a polypeptide, analog as described herein.

5

Thus, what is important for analogs, derivatives and fragments is that they possess at least a degree of the antigenicity/immunogenicity of the protein or polypeptide from which they are derived.

10

Also included are polypeptides which have fused thereto other compounds which alter the polypeptides biological or pharmacological properties i.e. polyethylene glycol (PEG) to increase half-life; leader or secretory amino acid sequences for ease of purification; prepro- and pro- sequences; and (poly)saccharides.

15

Furthermore, in those situations where amino acid regions are found to be polymorphic, it may be desirable to vary one or more particular amino acids to more effectively mimic the different epitopes of the different Streptococcus strains.

20

Moreover, the polypeptides of the present invention can be modified by terminal -NH<sub>2</sub> acylation (eg. by acetylation, or thioglycolic acid amidation, terminal carboxy amidation, e.g. with ammonia or methylamine) to provide stability, increased hydrophobicity for linking or binding to a support or other molecule.

25

Also contemplated are hetero and homo polypeptide multimers of the polypeptide fragments and analogues. These polymeric forms include, for example, one or more polypeptides that have been cross-linked with cross-linkers such as

30



avidin/biotin, gluteraldehyde or dimethylsuperimide. Such polymeric forms also include polypeptides containing two or more tandem or inverted contiguous sequences, produced from multicistronic mRNAs generated by recombinant DNA  
5 technology.

In a further embodiment, the present invention also relates to chimeric polypeptides which comprise one or more polypeptides or fragments or analogs thereof as defined in  
10 the figures of the present application.

In a further embodiment, the present invention also relates to chimeric polypeptides comprising two or more polypeptides having a sequence chosen from SEQ ID NO: 2, or fragments or  
15 analogs thereof; provided that the polypeptides are linked as to formed a chimeric polypeptide.

In a further embodiment, the present invention also relates to chimeric polypeptides comprising two or more polypeptides having a sequence chosen from SEQ ID NO: 2 provided that the polypeptides are linked as to formed a chimeric polypeptide.  
20

Preferably, a fragment, analog or derivative of a polypeptide of the invention will comprise at least one  
25 antigenic region i.e. at least one epitope.

In order to achieve the formation of antigenic polymers (i.e. synthetic multimers), polypeptides may be utilized having bishaloacetyl groups, nitroarylhalides, or the like,  
30 where the reagents being specific for thio groups.

Therefore, the link between two mercapto groups of the different polypeptides may be a single bond or may be composed of a linking group of at least two, typically at

least four, and not more than 16, but usually not more than about 14 carbon atoms.

In a particular embodiment, polypeptide fragments and  
5 analogs of the invention do not contain a starting residue, such as methionine (Met) or valine (Val). Preferably, polypeptides will not incorporate a leader or secretory sequence (signal sequence). The signal portion of a polypeptide of the invention may be determined according to  
10 established molecular biological techniques. In general, the polypeptide of interest may be isolated from a streptococcal culture and subsequently sequenced to determine the initial residue of the mature protein and therefore the sequence of the mature polypeptide.

15 It is understood that polypeptides can be produced and/or used without their start codon (methionine or valine) and/or without their leader peptide to favor production and purification of recombinant polypeptides. It is known that  
20 cloning genes without sequences encoding leader peptides will restrict the polypeptides to the cytoplasm of E. coli and will facilitate their recovery (Glick, B.R. and Pasternak, J.J. (1998) Manipulation of gene expression in prokaryotes. In "Molecular biotechnology: Principles and  
25 applications of recombinant DNA", 2nd edition, ASM Press, Washington DC, p.109-143).

According to another aspect of the invention, there are also provided (i) a composition of matter containing a  
30 polypeptide of the invention, together with a carrier, diluent or adjuvant; (ii) a pharmaceutical composition comprising a polypeptide of the invention and a carrier, diluent or adjuvant; (iii) a vaccine comprising a

polypeptide of the invention and a carrier, diluent or adjuvant; (iv) a method for inducing an immune response against Streptococcus, in a host, by administering to the host, an immunogenically effective amount of a polypeptide  
5 of the invention to elicit an immune response, e.g., a protective immune response to Streptococcus; and particularly, (v) a method for preventing and/or treating a Streptococcus infection, by administering a prophylactic or therapeutic amount of a polypeptide of the invention to a  
10 host in need.

According to another aspect of the invention, there are also provided (i) a composition of matter containing a polynucleotide of the invention, together with a carrier,  
15 diluent or adjuvant; (ii) a pharmaceutical composition comprising a polynucleotide of the invention and a carrier, diluent or adjuvant; (iii) a method for inducing an immune response against Streptococcus, in a host, by administering to the host, an immunogenically effective amount of a  
20 polynucleotide of the invention to elicit an immune response, e.g., a protective immune response to Streptococcus; and particularly, (iv) a method for preventing and/or treating a Streptococcus infection, by administering a prophylactic or therapeutic amount of a  
25 polynucleotide of the invention to a host in need.

Before immunization, the polypeptides of the invention can also be coupled or conjugated to carrier proteins such as tetanus toxin, diphtheria toxin, hepatitis B virus surface  
30 antigen, poliomyelitis virus VP1 antigen or any other viral or bacterial toxin or antigen or any suitable proteins to stimulate the development of a stronger immune response. This coupling or conjugation can be done chemically or

genetically. A more detailed description of peptide-carrier conjugation is available in Van Regenmortel, M.H.V., Briand J.P., Muller S., Plaué S., «Synthetic Polypeptides as antigens» in Laboratory Techniques in Biochemistry and Molecular Biology, Vol.19 (ed.) Burdou, R.H. & Van Knippenberg P.H. (1988), Elsevier New York.

According to another aspect, there are provided pharmaceutical compositions comprising one or more Streptococcal polypeptides of the invention in a mixture with a pharmaceutically acceptable adjuvant. Suitable adjuvants include (1) oil-in-water emulsion formulations such as MF59<sup>TM</sup>, SAF<sup>TM</sup>, Ribi<sup>TM</sup>; (2) Freund's complete or incomplete adjuvant; (3) salts i.e. AlK(SO<sub>4</sub>)<sub>2</sub>, AlNa(SO<sub>4</sub>)<sub>2</sub>, AlNH<sub>4</sub>(SO<sub>4</sub>)<sub>2</sub>, Al(OH)<sub>3</sub>, AlPO<sub>4</sub>, silica, kaolin; (4) saponin derivatives such as Stimulon<sup>TM</sup> or particles generated therefrom such as ISCOMs (immunostimulating complexes); (5) cytokines such as interleukins, interferons, macrophage colony stimulating factor (M-CSF), tumor necrosis factor (TNF); (6) other substances such as carbon polynucleotides i.e. poly IC and poly AU, detoxified cholera toxin (CTB) and E.coli heat labile toxin for induction of mucosal immunity. A more detailed description of adjuvant is available in a review by M.Z.I Khan et al. in Pharmaceutical Research, vol. 11, No. 1 (1994) pp2-11, and also in another review by Gupta et al., in Vaccine, Vol. 13, No. 14, pp1263-1276 (1995) and in WO 99/24578. Preferred adjuvants include Quila<sup>TM</sup>, QS21<sup>TM</sup>, Alhydrogel<sup>TM</sup> and Adjuphos<sup>TM</sup>.

Pharmaceutical compositions of the invention may be administered parenterally by injection, rapid infusion,

nasopharyngeal absorption, dermoabsorption, or buccal or oral.

Pharmaceutical compositions of the invention are used for  
5 the prophylaxis or treatment of streptococcal infection  
and/or diseases and symptoms mediated by streptococcal  
infection as described in Manual of Clinical Microbiology,  
P.R. Murray (Ed, in chief), E.J. Baron, M.A. Pfaller, F.C.  
Tenover and R.H. Tenover. ASM Press, Washington, D.C.  
10 seventh edition, 1999, 1773p. In one embodiment,  
pharmaceutical compositions of the present invention are  
used for the prophylaxis or treatment of pharyngitis,  
erysipelas and impetigo, scarlet fever, and invasive  
diseases such as bacteremia and necrotizing fasciitis and  
15 also toxic shock. In one embodiment, pharmaceutical  
compositions of the invention are used for the treatment or  
prophylaxis of Streptococcus infection and/or diseases and  
symptoms mediated by Streptococcus infection, in particular  
group B Streptococcus (GBS or S.agalactiae), group A  
20 Streptococcus (Streptococcus pyogenes), S.pneumoniae,  
S.dysgalactiae, S.uberis, S.nocardia as well as  
Staphylococcus aureus. In a further embodiment, the  
Streptococcus infection is group B Streptococcus (GBS or  
S.agalactiae).  
25  
In a further embodiment, the invention provides a method for  
prophylaxis or treatment of Streptococcus infection in a  
host susceptible to Streptococcus infection comprising  
administering to said host a prophylactic or therapeutic  
30 amount of a composition of the invention.

In a further embodiment, the invention provides a method for prophylaxis or treatment of GBS infection in a host susceptible to GBS infection comprising administering to said host a prophylactic or therapeutic amount of a composition of the invention.

As used in the present application, the term "host" includes mammals. In a further embodiment, the mammal is a member of a dairy herd. In a further embodiment, the mammal is an expectant mother. In a further embodiment, the mammal is human. In a further embodiment, the host is a pregnant woman. In a further embodiment, the host is a non-pregnant adult. In a further embodiment, the host is a neonate or an infant.

In a particular embodiment, pharmaceutical compositions are administered to those hosts at risk of streptococcus infection such as infants, elderly and immunocompromised hosts.

Pharmaceutical compositions are preferably in unit dosage form of about 0.001 to 100 µg/kg (antigen/body weight) and more preferably 0.01 to 10 µg/kg and most preferably 0.1 to 1 µg/kg 1 to 3 times with an interval of about 1 to 6 week intervals between immunizations.

Pharmaceutical compositions are preferably in unit dosage form of about 0.1 µg to 10 mg and more preferably 1µg to 1 mg and most preferably 10 to 100 µg 1 to 3 times with an interval of about 1 to 6 week intervals between immunizations.

According to another aspect, there are provided polynucleotides encoding polypeptides characterized by the amino acid sequence comprising SEQ ID NO: 2 or fragments or analogs thereof.

5

In one embodiment, polynucleotides are those illustrated in SEQ ID No: 1 which may include the open reading frames (ORF), encoding the polypeptides of the invention.

10 It will be appreciated that the polynucleotide sequences illustrated in the figures may be altered with degenerate codons yet still encode the polypeptides of the invention. Accordingly the present invention further provides polynucleotides which hybridize to the polynucleotide  
15 sequences herein above described (or the complement sequences thereof) having 80% identity between sequences. In one embodiment, at least 85% identity between sequences. In one embodiment, at least 90% identity between sequences. In a further embodiment, polynucleotides are hybridizable under  
20 stringent conditions i.e. having at least 95% identity. In a further embodiment, more than 97% identity.

Suitable stringent conditions for hybridation can be readily determined by one of skilled in the art (see for example  
25 Sambrook et al., (1989) Molecular cloning : A Laboratory Manual, 2<sup>nd</sup> ed, Cold Spring Harbor, N.Y.; Current Protocols in Molecular Biology, (1999) Edited by Ausubel F.M. et al., John Wiley & Sons, Inc., N.Y.).

30 In a further embodiment, the present invention provides polynucleotides that hybridize under stringent conditions to either

(a) a DNA sequence encoding a polypeptide or

(b) the complement of a DNA sequence encoding a polypeptide;  
wherein said polypeptide comprises SEQ ID NO:2 or fragments or analogs thereof.

5

In a further embodiment, the present invention provides polynucleotides that hybridize under stringent conditions to either

- (a) a DNA sequence encoding a polypeptide or
  - 10 (b) the complement of a DNA sequence encoding a polypeptide;
- wherein said polypeptide comprises SEQ ID NO: 2.

15 In a further embodiment, the present invention provides polynucleotides that hybridize under stringent conditions to either

- (a) a DNA sequence encoding a polypeptide or
  - (b) the complement of a DNA sequence encoding a polypeptide;
- 20 wherein said polypeptide comprises at least 10 contiguous amino acid residues from a polypeptide comprising SEQ ID NO:2 or fragments or analogs thereof.

In a further embodiment, the present invention provides  
25 polynucleotides that hybridize under stringent conditions to either

- (a) a DNA sequence encoding a polypeptide or
  - (b) the complement of a DNA sequence encoding a polypeptide;
- 30 wherein said polypeptide comprises at least 10 contiguous amino acid residues from a polypeptide comprising SEQ ID NO: 2.

In a further embodiment, polynucleotides of the invention  
35 are those encoding polypeptides of the invention illustrated in SEQ ID NO: 2 or fragments or analogs thereof.



In a further embodiment, polynucleotides of the invention are those illustrated in SEQ ID NO: 1 encoding polypeptides of the invention or fragments or analogs thereof.

5

In a further embodiment, polynucleotides of the invention are those encoding polypeptides of the invention illustrated in SEQ ID NO: 2.

10 In a further embodiment, polynucleotides of the invention are those illustrated in SEQ ID NO: 1 encoding polypeptides of the invention.

As will be readily appreciated by one skilled in the art,  
15 polynucleotides include both DNA and RNA.

The present invention also includes polynucleotides complementary to the polynucleotides described in the present application.

20

In a further aspect, polynucleotides encoding polypeptides of the invention, or fragments, analogs or derivatives thereof, may be used in a DNA immunization method. That is, they can be incorporated into a vector which is replicable  
25 and expressible upon injection thereby producing the antigenic polypeptide in vivo. For example polynucleotides may be incorporated into a plasmid vector under the control of the CMV promoter which is functional in eukaryotic cells. Preferably the vector is injected intramuscularly.

30

According to another aspect, there is provided a process for producing polypeptides of the invention by recombinant techniques by expressing a polynucleotide encoding said

polypeptide in a host cell and recovering the expressed polypeptide product.

Alternatively, the polypeptides can be produced according to established synthetic chemical techniques i.e. solution phase or solid phase synthesis of oligopeptides which are ligated to produce the full polypeptide (block ligation).

General methods for obtention and evaluation of polynucleotides and polypeptides are described in the following references: Sambrook et al, Molecular Cloning: A Laboratory Manual, 2nd ed, Cold Spring Harbor, N.Y., 1989; Current Protocols in Molecular Biology, Edited by Ausubel F.M. et al., John Wiley and Sons, Inc. New York; PCR Cloning Protocols, from Molecular Cloning to Genetic Engineering, Edited by White B.A., Humana Press, Totowa, New Jersey, 1997, 490 pages; Protein Purification, Principles and Practices, Scopes R.K., Springer-Verlag, New York, 3rd Edition, 1993, 380 pages; Current Protocols in Immunology, Edited by Coligan J.E. et al., John Wiley & Sons Inc., New York.

The present invention provides a process for producing a polypeptide comprising culturing a host cell of the invention under conditions suitable for expression of said polypeptide.

For recombinant production, host cells are transfected with vectors which encode the polypeptides of the invention, and then cultured in a nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying the genes. Suitable vectors are those that are

viable and replicable in the chosen host and include chromosomal, non-chromosomal and synthetic DNA sequences e.g. bacterial plasmids, phage DNA, baculovirus, yeast plasmids, vectors derived from combinations of plasmids and phage DNA. The polypeptide sequence may be incorporated in the vector at the appropriate site using restriction enzymes such that it is operably linked to an expression control region comprising a promoter, ribosome binding site (consensus region or Shine-Dalgarno sequence), and optionally an operator (control element). One can select individual components of the expression control region that are appropriate for a given host and vector according to established molecular biology principles (Sambrook et al, Molecular Cloning: A Laboratory Manual, 2nd ed, Cold Spring Harbor, N.Y., 1989; Current Protocols in Molecular Biology, Edited by Ausubel F.M. et al., John Wiley and Sons, Inc. New York). Suitable promoters include but are not limited to LTR or SV40 promoter, E.coli lac, tac or trp promoters and the phage lambda P<sub>L</sub> promoter. Vectors will preferably incorporate an origin of replication as well as selection markers i.e. ampicillin resistance gene. Suitable bacterial vectors include pET, pQE70, pQE60, pQE-9, pD10 phagescript, psiX174, pbluescript SK, pbsks, pNH8A, pNH16a, pNH18A, pNH46A, ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 and eukaryotic vectors pBlueBacIII, pWLNEO, pSV2CAT, pOG44, pXT1, pSG, pSVK3, pBPV, pMSG and pSVL. Host cells may be bacterial i.e. E.coli, Bacillus subtilis, Streptomyces; fungal i.e. Aspergillus niger, Aspergillus nidulins; yeast i.e. Saccharomyces or eukaryotic i.e. CHO, COS.

Upon expression of the polypeptide in culture, cells are typically harvested by centrifugation then disrupted by

physical or chemical means (if the expressed polypeptide is not secreted into the media) and the resulting crude extract retained to isolate the polypeptide of interest.

Purification of the polypeptide from culture media or lysate may be achieved by established techniques depending on the properties of the polypeptide i.e. using ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, hydroxylapatite chromatography and lectin chromatography. Final purification may be achieved using HPLC.

The polypeptides may be expressed with or without a leader or secretion sequence. In the former case the leader may be removed using post-translational processing (see US 4,431,739; US 4,425,437; and US 4,338,397) or be chemically removed subsequent to purifying the expressed polypeptide.

According to a further aspect, the streptococcal polypeptides of the invention may be used in a diagnostic test for Streptococcus infection, in particular group B Streptococcus infection. Several diagnostic methods are possible, for example detecting Streptococcus organism in a biological sample, the following procedure may be followed:

- a) obtaining a biological sample from a host;
- b) incubating an antibody or fragment thereof reactive with a polypeptide of the invention with the biological sample to form a mixture; and
- c) detecting specifically bound antibody or bound fragment in the mixture which indicates the presence of Streptococcus.

Alternatively, a method for the detection of antibody specific to a Streptococcus antigen in a biological sample containing or suspected of containing said antibody may be performed as follows:

- 5 a) obtaining a biological sample from a host;
- b) incubating one or more polypeptides of the invention or fragments thereof with the biological sample to form a mixture; and
- c) detecting specifically bound antigen or bound
- 10 fragment in the mixture which indicates the presence of antibody specific to Streptococcus.

One of skill in the art will recognize that this diagnostic test may take several forms, including an immunological test

15 such as an enzyme-linked immunosorbent assay (ELISA), a radioimmunoassay or a latex agglutination assay, essentially to determine whether antibodies specific for the protein are present in an organism.

20 The DNA sequences encoding polypeptides of the invention may also be used to design DNA probes for use in detecting the presence of Streptococcus in a biological sample suspected of containing such bacteria. The detection method of this invention comprises:

- 25 a) obtaining the biological sample from a host;
- b) incubating one or more DNA probes having a DNA sequence encoding a polypeptide of the invention or fragments thereof with the biological sample to form a mixture; and
- 30 c) detecting specifically bound DNA probe in the mixture which indicates the presence of Streptococcus bacteria.

The DNA probes of this invention may also be used for detecting circulating Streptococcus i.e. group B Streptococcus nucleic acids in a sample, for example using a polymerase chain reaction, as a method of diagnosing Streptococcus infections. The probe may be synthesized using conventional techniques and may be immobilized on a solid phase, or may be labelled with a detectable label. A preferred DNA probe for this application is an oligomer having a sequence complementary to at least about 6 contiguous nucleotides of the group B Streptococcus polypeptides of the invention. In a further embodiment, the preferred DNA probe will be an oligomer having a sequence complementary to at least about 15 contiguous nucleotides of the group B Streptococcus polypeptides of the invention. In a further embodiment, the preferred DNA probe will be an oligomer having a sequence complementary to at least about 30 contiguous nucleotides of the group B Streptococcus polypeptides of the invention. In a further embodiment, the preferred DNA probe will be an oligomer having a sequence complementary to at least about 50 contiguous nucleotides of the group B Streptococcus polypeptides of the invention.

Another diagnostic method for the detection of Streptococcus in a host comprises:

- a) labelling an antibody reactive with a polypeptide of the invention or fragment thereof with a detectable label;
- b) administering the labelled antibody or labelled fragment to the host; and

- c) detecting specifically bound labelled antibody or labelled fragment in the host which indicates the presence of Streptococcus.

5 Alternatively, a method for the detection of antibody specific to a Streptococcus antigen in a biological sample containing or suspected of containing said antibody may be performed as follows:

- a) obtaining a biological sample from a host;
- 10 b) incubating one or more Streptococcus polypeptides of the invention or fragments thereof with the biological sample to form a mixture; and
- c) detecting specifically bound antigen or bound fragment in the mixture which indicates the
- 15 presence of antibody specific to Streptococcus.

One of skill in the art will recognize that the diagnostic test may take several forms, including an immunological test such as an enzyme-linked immunosorbent assay (ELISA), a

20 radioimmunoassay or a latex agglutination assay, essentially to determine whether antibodies specific for the protein are present in an organism.

The DNA sequences encoding polypeptides of the invention may also be used to design DNA probes for use in detecting the

25 presence of Streptococcus in a biological sample suspected of containing such bacteria. The detection method of this invention comprises:

- a) obtaining the biological sample from a host;
- 30 b) incubating one or more DNA probes having a DNA sequence encoding a polypeptide of the invention or fragments thereof with the biological sample to form a mixture; and

- c) detecting specifically bound DNA probe in the mixture which indicates the presence of Streptococcus bacteria.

5 According to one aspect, the present invention provides the use of an antibody for treatment and/or prophylaxis of streptococcal infections.

A further aspect of the invention is the use of the  
10 Streptococcus polypeptides of the invention as immunogens for the production of specific antibodies for the diagnosis and in particular the treatment of streptococcus infection. Suitable antibodies may be determined using appropriate screening methods, for example by measuring the ability of a  
15 particular antibody to passively protect against streptococcus infection in a test model. One example of an animal model is the mouse model described in the examples herein. The antibody may be a whole antibody or an antigen-binding fragment thereof and may belong to any  
20 immunoglobulin class. The antibody or fragment may be of animal origin, specifically of mammalian origin and more specifically of murine, rat or human origin. It may be a natural antibody or a fragment thereof, or if desired, a recombinant antibody or antibody fragment. The term  
25 recombinant antibody or antibody fragment means antibody or antibody fragment which was produced using molecular biology techniques. The antibody or antibody fragments may be polyclonal, or preferably monoclonal. It may be specific for a number of epitopes associated with the Group B  
30 Streptococcus polypeptides but is preferably specific for one.



A further aspect of the invention is the use of the antibodies directed to the polypeptides of the invention for passive immunization. One could use the antibodies described in the present application.

5

A further aspect of the invention is a method for immunization, whereby an antibody raised by a polypeptide of the invention is administered to a host in an amount sufficient to provide a passive immunization.

10

In a further embodiment, the invention provides the use of a pharmaceutical composition of the invention in the manufacture of a medicament for the prophylactic or therapeutic treatment of streptococcal infection.

15

In a further embodiment, the invention provides a kit comprising a polypeptide of the invention for detection or diagnosis of streptococcal infection.

20 Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. All publications, patent applications, patents, and other references mentioned herein are incorporated by  
25 reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

30

#### EXAMPLE 1

This example illustrates the identification of Group B streptococcal BVH-A4 gene.

Chromosomal DNA was isolated from different Group B streptococcal strains as previously described (Jayarao BM et al. 1991. J. Clin. Microbiol. 29:2774-2778). A  $\lambda$ ZAPExpress genomic library was constructed using chromosomal DNA

5 purified from the serotype III Group B streptococcal strain COH1 (Children's Hospital and Medical Center, Seattle, WA, USA) and screened according to the manufacturer's instructions (Stratagene, La Jolla, CA) with a pool of human normal sera. Briefly, the purified chromosomal DNA was

10 partially digested with *tsp509I* restriction enzyme, and the resulting fragments were electrophoresed on a 1% agarose gel (Bio-Rad). Fragments in the 5- to 10-kb size range were extracted from the gel and ligated to the *EcoRI* arms of

15  $\lambda$ ZAPExpress vector and the vector was encapsidated using the Gigapack II packaging extract (Stratagene). The recombinant phages were used to infect *E. coli* XL1-Blue MRF' [ $\Delta(mcrA)183\Delta(mcrCB-hsdSMR-mrr)173\ endA1\ supE44\ thi-1\ recA1\ gyrA96\ relA1\ lac\ (F'\ proAB\ lacI^qZAM15\ Tn10\ [Tet^R])$ ], which was then plated onto LB agar. The resulting plaques

20 were lifted onto Hybond-C nitrocellulose membranes (Amersham Pharmacia Biotech, Baie d'Urfée, Canada) pre-impregnated with 10 mM Isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG: ICN Biomedicals Inc., Costa Mesa, CA). The membranes were blocked using phosphate-buffered saline (PBS) with 3% skim

25 milk and were sequentially incubated with the pooled of human sera, peroxidase-labeled goat anti-human immunoglobulins antisera (Jackson Immunoresearch Laboratories Inc., West Grove, PA) and substrate. Positive plaques were isolated, purified twice, and the recombinant

30 pBK-CMV plasmids (Stratagene) were excised with the ExAssist helper phage (Stratagene) according to the manufacturer's instructions. Immunoblots using phagemid vectors containing the cloned insert revealed that the pooled human sera reacted with a protein band with an approximate molecular

35 weight of 100 kDa for the clone A1. This clone was then identified as BVH-A4. The sequence of the insert was determined using the TAQ Dye Deoxy Terminator Cycle

Sequencing Kit with an Applied Biosystems Inc. (Foster City, CA) automated sequencer model 373A according to the manufacturer's recommendations.

5

## EXAMPLE 2

This example illustrates the cloning of Group B streptococcal BVH-A4 gene.

10 The coding region of Group B streptococcal BVH-A4 (SEQ ID NO: 1) gene without the region coding for the leader peptide was amplified by PCR (DNA Thermal Cycler GeneAmp PCR system 2400 Perkin Elmer, San Jose, CA) from genomic DNA of serotype III Group B streptococcal strain COH1 using  
15 oligonucleotide primers that contained base extensions for the addition of restriction sites *Nco*I (CCATGG) and *Not*I (GCGGCCGC). The oligonucleotide primers (Table 1) DMAR800 and OCRR588 were used to amplify the BVH-A4 gene. PCR products were purified from agarose gel using a QIAquick gel  
20 extraction kit from QIAGEN following the manufacturer's instructions (Chatsworth, CA), and digested with *Nco*I and *Not*I (Pharmacia Canada Inc, Baie d'Urfée, Canada). The pET-21d(+) vector (Novagen, Madison, WI) was digested with *Nco*I and *Not*I and purified from agarose gel using a QIAquick gel  
25 extraction kit from QIAGEN (Chatsworth, CA). The *Nco*I-*Not*I PCR product was ligated to the *Nco*I-*Not*I pET-21d(+) expression vector. The ligated product was transformed into *E. coli* strain DH5 $\alpha$  [ $\phi$ 80dlacZAM15  $\Delta$ (lacZYA-argF)U169 *endA*1 *recA*1 *hsdR*17(*r<sub>K</sub>-m<sub>K</sub>*+) *deoR* *thi*-1 *supE*44  $\lambda$ <sup>-</sup>  
30 *gyrA*96 *relA*1] (Gibco BRL, Gaithersburg, MD) according to the method of Simanis (Hanahan, D. DNA Cloning, 1985, D.M. Glover (ed), pp. 109-135). Recombinant pET-21d(+) plasmid (rpET21d(+)) containing BVH-A4 gene was purified using a QIAGEN plasmid kit (Chatsworth, CA) and DNA insert was  
35 sequenced (Taq Dye Deoxy Terminator Cycle Sequencing kit, ABI, Foster City, CA).

It was determined that the open reading frame (ORF) which codes for BVH-A4 gene (SEQ ID NO: 1) contains 3168-bp and encodes a 1055 amino acid residues polypeptide with a predicted pI of 7.97 and a predicted molecular mass of 118,151.59 Da. Analysis of the predicted amino acid residues sequence (SEQ ID NO :2) using the Spscan software (Wisconsin Sequence Analysis Package; Genetics Computer Group) suggested the existence of a 22 amino acid residues signal peptide (MTKKHLKTLALALTTSVVTYS), which ends with a cleavage site situated between a serine and a glutamine residues.

Table 1. Oligonucleotide primers used for PCR amplifications of Group B streptococcal BVH-A4 gene

Gene	Primers I.D.	Restriction site	Vector	Sequence
BVH-A4	DMAR800	<i>Nco</i> I	pET21d	5'-GCGCCCATGGTGCAGGAGGTATATGGT TAGAAAG-3' (SEQ ID No: 3)
BVH-A4	OCRR587	<i>Bam</i> HI	pET21b	5'-GGTGGATCCGAGAAAGGCTTTATTGTA ATG-3' (SEQ ID No: 4)
BVH-A4	OCRR588	<i>Not</i> I	pET21d or pET21b	5'-CATATTAATTGCGGCCGCTTTTCTTGC TCGTTTCC-3' (SEQ ID No: 5)
BVH-A4	DMAR752	<i>Bam</i> HI	pCMV-GH	5'-CGTTGGATCCTCAGGAGGTATATGGAT TAGAAAG-3' (SEQ ID No: 6)
BVH-A4	DMAR753	<i>Sal</i> I	pCMV-GH	5'-CATCGTCGACTTATTTTCTTGCTCGTT TTCC-3' (SEQ ID No: 7)

### EXAMPLE 3

This example describes the PCR amplification of Group B streptococcal BVH-A4 gene from other Group B strains

To confirm the presence by PCR amplification of BVH-A4 (SEQ ID NO :1) gene, the following 11 serologically distinct Group B streptococcal strains were used: C388/90 (serotype Ia/c), ATCC12401 (serotype Ib), ATCC27591 (serotype Ic), NCS246 (serotype II/R), NCS954 (serotype III), NCS97SR331 (serotype IV), NCS535 (serotype V), NCS9842 (serotype VI),

NCS7271 (serotype VII), NCS970886 (serotype VIII), ATCC27956 (bovine isolate). These strains were obtained from the American Type Culture Collection (Rockville, MD, USA) and National Centre for Streptococcus, Provincial Laboratory of Public Health for Northern Alberta (Edmonton, Canada). The E.coli strain XL1-Blue MRF' was used in these experiments as negative control. Chromosomal DNA was isolated from each Group B streptococcal strain as previously described (Jayarao BM et al. 1991. J. Clin. Microbiol. 29:2774-2778).

BVH-A4 (SEQ ID NO :1) gene was amplified by PCR (DNA Thermal Cycler GeneAmp PCR system 2400 Perkin Elmer, San Jose, CA) from the genomic DNA purified from the 11 Group B streptococcal strains, and the control E.coli strain using the oligonucleotides presented in Table 1. The oligonucleotide primers OCRR587 and OCRR588 were used to amplify the BVH-A4 (SEQ ID NO :1) gene. PCR was performed with 35 cycles of 30 sec at 94°C, 30 sec at 55°C and 150 sec at 72°C and a final elongation period of 30 min at 72°C. The PCR products were size fractionated in 1% agarose gels and were visualized by ethidium bromide staining. The results of these PCR amplifications are presented in Table 2. The analysis of the amplification products revealed that BVH-A4 (SEQ ID NO :1) gene was present in the genome of all of the 11 Group B streptococcal strains tested. No such product was detected when the control E.coli DNA was submitted to identical PCR amplifications with these oligonucleotide primers.

Table 2. Identification of BVH-A4 gene by PCR amplification

Group B streptococcal isolates	Strains identification
	<u>BVH-A4</u>
C388/90 (serotype Ia/c)	+
ATCC12401 (serotype Ib)	+
ATCC27591 (serotype Ic)	+
NCS246 (serotype II/R)	+
NCS954 (serotype III)	+
NCS97SR331 (serotype IV)	+

NCS535 (serotype V)	+
NCS9842 (serotype VI)	+
NCS7271 (serotype VII)	+
NCS970886 (serotype VIII)	+
ATCC27956 (bovine isolate)	+
<u>E.coli</u> control strain XL1 Blue MRF'	-

**EXAMPLE 4**

This example illustrates the cloning of Group B streptococcal BVH-A4 gene in CMV plasmid pCMV-GH.

The DNA coding region of Group B streptococcal BHV-A4 (SEQ ID NO :1) without the leader peptide was inserted in phase downstream of a human growth hormone (hGH) gene which was under the transcriptional control of the cytomegalovirus (CMV) promotor in the plasmid vector pCMV-GH (Tang et al., Nature, 1992, 356 :152). The CMV promotor is non functional plasmid in E.coli cells but active upon administration of the plasmid in eukaryotic cells. The vector also incorporated the ampicillin resistance gene.

The coding regions of BVH-A4 (SEQ ID NO: 1) gene without its leader peptide regions was amplified by PCR (DNA Thermal Cycler GeneAmp PCR system 2400 Perkin Elmer, San Jose, CA) from genomic DNA of serotype III Group B streptococcal strain COH1 using oligonucleotide primers that contained base extensions for the addition of restriction sites *Bam*HI (GGATCC) and *Sal*I (GTCGAC). The oligonucleotide primers DMAR752 and DMAR753 were used to amplify the BVH-A4 (SEQ ID NO :1) gene. The PCR product was purified from agarose gel using a QIAquick gel extraction kit from QIAgen (Chatsworth, CA), digested with restriction enzymes (Pharmacia Canada Inc, Baie d'Urfée, Canada). The pCMV-GH vector (Laboratory of Dr. Stephen A. Johnston, Department of Biochemistry, The University of Texas, Dallas, Texas) was digested with *Bam*HI and *Sal*I and purified from agarose gel using the QIAquick

gel extraction kit from QIAGEN (Chatsworth, CA). The *Bam*HI-*S*alI DNA fragments were ligated to the *Bam*HI-*S*alI pCMV-GH vector to create the hGH-BVH-A4 fusion protein under the control of the CMV promoter. The ligated product was transformed into *E. coli* strain DH5 $\alpha$  [ $\phi$ 80dlacZ $\Delta$ M15  $\Delta$ (lacZYA-argF)U169 *endA*1 *recA*1 *hsdR*17(*r*<sub>K</sub>-*m*<sub>K</sub>+) *deoR* *thi*-1 *supE*44  $\lambda$ <sup>-</sup> *gyrA*96 *relA*1] (Gibco BRL, Gaithersburg, MD) according to the method of Simanlis (Hanahan, D. DNA Cloning, 1985, D.M. Glover (ed), pp. 109-135). The recombinant pCMV plasmid was purified using a QIAGEN plasmid kit (Chatsworth, CA) and the nucleotide sequence of the DNA insert was verified by DNA sequencing.

#### 15 EXAMPLE 5

This example illustrates the use of DNA to elicit an immune response to Group B streptococcal BVH-A4 protein antigen.

Groups of 8 female BALB/c mice (Charles River, St-Constant, Québec, Canada) were immunized by intramuscular injection of 100  $\mu$ l three times at two- or three-week intervals with 50  $\mu$ g of recombinant pCMV-GH encoding BVH-A4 (SEQ ID NO :1) gene in presence of 50  $\mu$ g of granulocyte-macrophage colony-stimulating factor (GM-CSF)- expressing plasmid pCMV-GH-GM-CSF (Laboratory of Dr. Stephen A. Johnston, Department of Biochemistry, The University of Texas, Dallas, Texas). As control, groups of mice were injected with 50  $\mu$ g of pCMV-GH in presence of 50  $\mu$ g of pCMV-GH-GM-CSF. Blood samples were collected from the orbital sinus prior to each immunization and seven days following the third injection and serum antibody responses were determined by ELISA using purified BVH-A4-His•Tag recombinant proteins as coating antigen.

#### 35 EXAMPLE 6

This example illustrates the production and purification of recombinant Group B streptococcal BVH-A4 protein.

The recombinant pET-21d(+) plasmid with BVH-A4 gene  
5 corresponding to the SEQ ID NO: 1 was used to transform by electroporation (Gene Pulser II apparatus, BIO-RAD Labs, Mississauga, Canada) E.coli strain BL21(DE3) (F<sup>+</sup>ompT hsdS<sub>B</sub> (r<sup>-</sup><sub>m<sub>B</sub></sub>) gal dcm (DE3)) (Novagen, Madison, WI). In this strain of E.coli, the T7 promotor controlling expression of the  
10 recombinant protein is specifically recognized by the T7 RNA polymerase (present on the  $\lambda$ DE3 prophage) whose gene is under the control of the lac promotor which is inducible by isopropyl- $\beta$ -d-thio-galactopyranoside (IPTG). The transformants BL21(DE3)/rpET were grown at 37°C with  
15 agitation at 250 rpm in LB broth (peptone 10g/L, yeast extract 5g/L, NaCl 10g/L) containing 100  $\mu$ g of carbenicillin (Sigma-Aldrich Canada Ltd., Oakville, Canada) per ml until the A<sub>600</sub> reached a value of 0.6. In order to induce the production of Group B streptococcal BVH-A4-His•Tag  
20 recombinant protein, the cells were incubated for 3 additional hours in the presence of IPTG at a final concentration of 1 mM. Induced cells from a 500 ml culture were pelleted by centrifugation and frozen at -70°C.

25 The purification of the recombinant proteins from the soluble cytoplasmic fraction of IPTG-induced BL21(DE3)/rpET21d(+) was done by affinity chromatography based on the properties of the His•Tag sequence (6 consecutive histidine residues) to bind to divalent cations  
30 (Ni<sup>2+</sup>) immobilized on the His•Bind metal chelation resin. Briefly, the pelleted cells obtained from a 500 mL culture induced with IPTG was resuspended in lysis buffer (20 mM Tris, 500 mM NaCl, 10 mM imidazole, pH 7.9) containing 1 mM PMSF, sonicated and centrifuged at 12,000 X g for 20 min to  
35 remove debris. The supernatant was deposited on a Ni-NTA agarose column (Qiagen, Mississauga, Ontario, Canada). The Group B streptococcal BVH-A4-His•Tag recombinant protein was



eluted with 250 mM imidazole-500 mM NaCl-20 mM Tris pH 7.9. The removal of the salt and imidazole from the samples was done by dialysis against PBS at 4°C. The quantities of recombinant proteins obtained from the soluble fraction of *E. coli* were estimated by MicroBCA (Pierce, Rockford, Illinois).

#### EXAMPLE 7

10 This example illustrates the accessibility to antibodies of the Group B streptococcal BVH-A4 protein at the surface of Group B streptococcal strains.

Bacteria were grown in Todd Hewitt (TH) broth (Difco Laboratories, Detroit MI) with 0.5% Yeast extract (Difco Laboratories) and 0.5% peptone extract (Merck, Darmstadt, Germany) at 37°C in a 8% CO<sub>2</sub> atmosphere to give an OD<sub>490nm</sub> of 0.600 (~10<sup>8</sup> CFU/ml). Dilutions of anti-BVH-A4 or control sera were then added and allowed to bind to the cells, which were  
20 incubated for 2 h at 4°C. Samples were washed 4 times in blocking buffer [phosphate-buffered saline (PBS) containing 2% bovine serum albumin (BSA)], and then 1 ml of goat fluorescein (FITC)-conjugated anti-mouse IgG + IgM diluted in blocking buffer was added. After an additional incubation  
25 of 60 min at room temperature, samples were washed 4 times in blocking buffer and fixed with 0.25 % formaldehyde in PBS buffer for 18-24 h at 4°C. Cells were washed 2 times in PBS buffer and resuspended in 500 µl of PBS buffer. Cells were kept in the dark at 4°C until analyzed by flow cytometry  
30 (Epics® XL; Beckman Coulter, Inc.).

#### EXAMPLE 8

35 This example illustrates the protection against fatal Group B streptococcal infection induced by passive immunization of mice with rabbit hyper-immune sera.

New Zealand rabbits (Charles River laboratories, St-Constant, Canada) were injected subcutaneously at multiple sites with 50  $\mu$ g and 100  $\mu$ g of BVH-A4-His•Tag protein that was produced and purified as described in Example 6 and adsorbed to Alhydrogel adjuvant (Superfos Biosector a/s). Rabbits were immunized three times at three-week intervals with the BVH-A4-His•Tag protein. Blood samples were collected three weeks after the third injection. The antibodies present in the serum were purified by precipitation using 40% saturated ammonium sulfate. Groups of 10 female CD-1 mice (Charles River) were injected intravenously with 500  $\mu$ l of purified serum collected either from BVH-A4-His•Tag immunized rabbits, or rabbits immunized with an unrelated control recombinant protein. Eighteen hours later the mice were challenged with approximately  $8 \times 10^4$  CFU of the Group B streptococcal strain C388/90 (Ia/c). Samples of the Group B streptococcal challenge inoculum were plated on blood agar plates to determine the CFU and to verify the challenge dose. Deaths were recorded for a period of 14 days.

#### EXAMPLE 9

This example illustrates the protection of mice against fatal Group B streptococcal infection induced by immunization.

Groups of 8 female CD-1 mice (Charles River) were immunized subcutaneously three times at three-week intervals with 20  $\mu$ g of either BVH-A4-His•Tag protein that was produced and purified as described in Example 6 in presence of 10  $\mu$ g of QuilA adjuvant (Cedarlane Laboratories Ltd, Hornby, Canada). The control mice were injected with QuilA adjuvant alone in PBS. Blood samples were collected from the orbital sinus on day 1, 22 and 43 prior to each immunization and seven days (day 50) following the third injection. Two weeks later the mice were challenged with approximately  $8 \times 10^4$  CFU of the

Group B streptococcal strain C388/90 (Ia/c). Samples of the Group B streptococcal challenge inoculum were plated on blood agar plates to determine the CFU and to verify the challenge dose. Deaths were recorded for a period of 14  
5 days.

**CLAIMS:**

1. An isolated polynucleotide comprising a polynucleotide chosen from:

5 (a) a polynucleotide encoding a polypeptide having at least 80% identity to a second polypeptide comprising a sequence chosen from: SEQ ID NO: 2 or fragments or analogs thereof;

10 (b) a polynucleotide encoding a polypeptide having at least 95% identity to a second polypeptide comprising a sequence chosen from: SEQ ID NO: 2 or fragments or analogs thereof;

(c) a polynucleotide encoding a polypeptide comprising a sequence chosen from: SEQ ID NO: 2 or fragments  
15 or analogs thereof;

(d) a polynucleotide encoding a polypeptide capable of raising antibodies having binding specificity for a polypeptide comprising a sequence chosen from: SEQ ID NO: 2 or fragments or analogs thereof;

20 (e) a polynucleotide encoding an epitope bearing portion of a polypeptide comprising a sequence chosen from SEQ ID NO: 2 or fragments or analogs thereof;

(f) a polynucleotide comprising a sequence chosen from SEQ ID NO: 1 or fragments or analogs thereof;

25 (g) a polynucleotide that is complementary to a polynucleotide in (a), (b), (c), (d), (e) or (f).

2. An isolated polynucleotide comprising a polynucleotide chosen from:

30 (a) a polynucleotide encoding a polypeptide having at least 80% identity to a second polypeptide comprising a sequence chosen from: SEQ ID NO: 2;

(b) a polynucleotide encoding a polypeptide having at least 95% identity to a second polypeptide comprising a sequence chosen from: SEQ ID NO: 2;

(c) a polynucleotide encoding a polypeptide  
5 comprising a sequence chosen from: SEQ ID NO: 2;

(d) a polynucleotide encoding a polypeptide capable of raising antibodies having binding specificity for a polypeptide comprising a sequence chosen from: SEQ ID NO: 2;

10 (e) a polynucleotide encoding an epitope bearing portion of a polypeptide comprising a sequence chosen from SEQ ID NO: 2;

(f) a polynucleotide comprising a sequence chosen from SEQ ID NO: 1;

15 (g) a polynucleotide that is complementary to a polynucleotide in (a), (b), (c), (d), (e) or (f).

3. The polynucleotide of claim 1, wherein said polynucleotide is DNA.

20

4. The polynucleotide of claim 2, wherein said polynucleotide is DNA.

5. The polynucleotide of claim 1, wherein said  
25 polynucleotide is RNA.

6. The polynucleotide of claim 2, wherein said polynucleotide is RNA.

30 7. An isolated polynucleotide that hybridizes under stringent conditions to either

(a) a DNA sequence encoding a polypeptide or  
(b) the complement of a DNA sequence encoding a polypeptide;

wherein said polypeptide comprises SEQ ID NO:2 or fragments or analogs thereof.

8. The polynucleotide of claim 1 that hybridizes under  
5 stringent conditions to either

- (a) a DNA sequence encoding a polypeptide or
- (b) the complement of a DNA sequence encoding a

polypeptide;

wherein said polypeptide comprises SEQ ID NO: 2, or  
10 fragments or analogs thereof.

9. The polynucleotide of claim 2 that hybridizes under stringent conditions to either

- (a) a DNA sequence encoding a polypeptide or
- 15 (b) the complement of a DNA sequence encoding a

polypeptide;

wherein said polypeptide comprises SEQ ID NO: 2.

10. The polynucleotide of claim 1 that hybridizes  
20 under stringent conditions to either

- (a) a DNA sequence encoding a polypeptide or
- (b) the complement of a DNA sequence encoding

a polypeptide;

wherein said polypeptide comprises at least 10 contiguous  
25 amino acid residues from a polypeptide comprising SEQ ID NO:  
2, or fragments or analogs thereof.

11. The polynucleotide of claim 2 that hybridizes under stringent conditions to either

- 30 (a) a DNA sequence encoding a polypeptide or
- (b) the complement of a DNA sequence encoding

a polypeptide;

wherein said polypeptide comprises at least 10 contiguous  
amino acid residues from a polypeptide comprising SEQ ID  
35 NO: 2.

12. A vector comprising the polynucleotide of claim 1, wherein said DNA is operably linked to an expression control region.

5 13. A vector comprising the polynucleotide of claim 2, wherein said DNA is operably linked to an expression control region.

14. A host cell transfected with the vector of  
10 claim 12.

15. A host cell transfected with the vector of claim 13.

15 16. A process for producing a polypeptide comprising culturing a host cell according to claim 14 under conditions suitable for expression of said polypeptide.

17. A process for producing a polypeptide comprising  
20 culturing a host cell according to claim 15 under condition suitable for expression of said polypeptide.

18. An isolated polypeptide comprising a polypeptide chosen from:

25 (a) a polypeptide having at least 80% identity to a second polypeptide having an amino acid sequence comprising: SEQ ID NO: 2, or fragments or analogs thereof;

(b) a polypeptide having at least 95% identity to a second polypeptide having an amino acid sequence  
30 comprising: SEQ ID NO: 2, or fragments or analogs thereof;

(c) a polypeptide comprising a sequence chosen from SEQ ID NO: 2, or fragments or analogs thereof;

(d) a polypeptide capable of raising antibodies having binding specificity for a polypeptide comprising SEQ ID NO: 2, or fragments or analogs thereof;

(e) an epitope bearing portion of a polypeptide  
5 comprising SEQ ID NO: 2, or fragments or analogs thereof;

(f) the polypeptide of (a), (b), (c), (d), or (e) wherein the N-terminal Met residue is deleted;

(g) the polypeptide of (a), (b), (c), (d), (e), or (f) wherein the secretory amino acid sequence is deleted.

10

19. An isolated polypeptide comprising a polypeptide chosen from:

(a) a polypeptide having at least 80% identity to a second polypeptide having an amino acid sequence  
15 comprising: SEQ ID NO: 2;

(b) a polypeptide having at least 95% identity to a second polypeptide having an amino acid sequence comprising: SEQ ID NO: 2;

(c) a polypeptide comprising a sequence chosen  
20 from SEQ ID NO: 2;

(d) a polypeptide capable of raising antibodies having binding specificity for a polypeptide comprising SEQ ID NO: 2;

(e) an epitope bearing portion of a polypeptide  
25 comprising SEQ ID NO: 2;

(f) the polypeptide of (a), (b), (c), (d), or (e) wherein the N-terminal Met residue is deleted;

(g) the polypeptide of (a), (b), (c), (d), (e), or (f) wherein the secretory amino acid sequence is deleted.

30

20. A chimeric polypeptide comprising two or more polypeptides having a sequence chosen from SEQ ID NO: 2, or fragments or analogs thereof; provided that the polypeptides are linked as to formed a chimeric polypeptide.



21. A chimeric polypeptide comprising two or more polypeptides having a sequence chosen from SEQ ID NO: 2; provided that the polypeptides are linked as to formed a chimeric polypeptide.

5

22. A pharmaceutical composition comprising a polypeptide according to any one of claims 18 to 21 and a pharmaceutically acceptable carrier, diluent or adjuvant.

10 23. A method for prophylactic or therapeutic treatment of GBS bacterial infection in a host susceptible to GBS infection comprising administering to said host a prophylactic or therapeutic amount of a composition according to claim 22.

15

24. A method according to claim 23 wherein the host is a neonate or an infant.

25. A method according to claim 24 wherein said  
20 infection causes sepsis, meningitis, pneumonia, cellulitis, osteomyelitis, septic arthritis, endocarditis or epiglottitis.

26. A method according to claim 23 wherein the host is a pregnant woman.

25

27. A method according to claim 26 wherein said  
infection causes mild urinary tract infection to life-  
threatening sepsis and meningitis, including also  
osteomyelitis, endocarditis, amniotitis, endometritis, wound  
30 infections (postcesarean and postepisiotomy), cellulitis or fasciitis.

28. A method according to claim 23 wherein the host is a non-pregnant adult.

29. A method according to claim 28 wherein said  
5 infection causes primary bacteremia, skin or soft tissue infection, pneumonia, urosepsis, endocarditis, peritonitis, meningitis or empyema.

30. A method according to claim 23 wherein the host is  
10 a member of a dairy herd.

31. A method according to claim 30 wherein said infection causes mastitis.

15 32. A method for diagnostic of GBS bacterial infection in an host susceptible to GBS infection comprising  
(a) obtaining a biological sample from a host;  
(b) incubating an antibody or fragment thereof reactive with a polypeptide according to any one of claims  
20 18 to 21 with the biological sample to form a mixture; and  
(c) detecting specifically bound antibody or bound fragment in the mixture which indicates the presence of Streptococcus.

25 33. A method for diagnostic of GBS bacterial infection in an host susceptible to GBS infection comprising  
(a) obtaining a biological sample from a host;  
(b) incubating one or more polypeptides according to any one of claims 18 to 21 or fragments thereof with the  
30 biological sample to form a mixture; and  
(c) detecting specifically bound antigen or bound fragment in the mixture which indicates the presence of antibody specific to Streptococcus.

34. Use of the pharmaceutical composition according to claim 22 in the manufacture of a medicament for the prophylactic or therapeutic treatment of streptococcal infection.

5

35. Kit comprising a polypeptide according to any one of claims 18 to 21 for detection or diagnosis of streptococcal infection.

Figure 1 (SEQ ID NO: 1)

1 ATGACAAAAA AACATCTTAA AACGCTTGCC TTGGCACTTA CTACAGTATC  
AGTAGTGACA  
 61 TACAGCCAGG AGGTATATGG ATTAGAAAGA GAGGAATCGG TCAAACAAGA  
 ACAAACCCAG  
 121 TCAGCTTCAG AAGATGATTG GTTCGAAGAA GATAATGAGA GGAAAACAAA  
 TGTTTCTAAA  
 181 GAGAATTCTA CTGTTGATGA AACAGTTAGT GATTTATTTT CTGATGGAAA  
 TAGTAATAAC  
 241 TCTAGTTCTA AAACCGAGTC AGTGGTAAGT GACCCTAAAC AAGTCCCCAA  
 AGCAAAACCA  
 301 GAGGTTACAC AAGAAGCAAG CAATTCTAGT AATGATGCTA GCAAAGTAGA  
 AGTACCAAAA  
 361 CAGGATACAG CTTCAAAAAA GGAAACTCTA GAAACATCAA CTTGGGAGGC  
 AAAAGATTTT  
 421 GTAAC TAGAG GGGATACTTT AGTAGGTTTT TCAAAATCTG GAATTAATAA  
 GTTATCTCAA  
 481 ACATCACACT TGGTTTTACC AAGTCATGCA GCAGATGGAA CTCAATTGAC  
 ACAAGTAGCT  
 541 AGCTTTGCTT TTA CTCCAGA TAAAAGACG GCCATTGCAG AATATACAAG  
 TAGGCTAGGA  
 601 GAAAATGGGA AACCGAGTCG TTTAGATATT GATCAGAAGG AAATTATTGA  
 TGAGGGAGAA  
 661 ATATTTAATG CTTACCAGTT GACTAAGCTT ACTATTCCAA ATGGTTATAA  
 GTCTATTGGT  
 721 CAAGATGCTT TTGTGGACAA TAAGAATATT GCTGAGGTTA ACCTTCCTGA  
 GAGTCTCGAG  
 781 ACTATTTTCTAG ACTATGCTTT TGCTCACATG TCTTTAAAC AAGTAAAGTT  
 ACCAGATAAC  
 841 CTAAAGGTCA TTGGAGAATT AGCTTTTTTTT GATAATCAGA TTGGTGTTAA  
 GCTTTACTTG  
 901 CCACGTCCTT TGATAAAATT AGCAGAACGC GCTTTCAAAT CTAATCGTAT  
 TCAAACAGTT  
 961 GAATTTTTTGG GAAGTAAGCT TAAGGTTATA GGAGAAGCAA GTTTTCAAGA  
 TAATAATCTG  
 1021 AGGAATGTGA TGCTTCCGGA TGGACTTGAA AAAATAGAAT CAGAAGCTTT  
 TACAGGAAAT  
 1081 CCAGGAGATG AACATTACAA CAATCAGGTT GTATTGCGCA CAAGGACAGG  
 CCAAAATCCA  
 1141 CATCAACTTG CGACTGAGAA TACTTACGTC AATCCGGACA AATCATTGTG  
 GCGTGCAACA  
 1201 CCTGATATGG ATTATACCAA ATGGTTAGAG GAAGATTTTA CCTATCAAAA  
 AAATAGTGTT  
 1261 ACAGGTTTTT CAAATAAAGG CTTACAAAAG GTAAGACGTA ATAAAACTT  
 AGAAATTCCA  
 1321 AAACAACACA ATGGTATTAC TATTACTGAA ATTGGTGATA ACGCTTTTCG  
 CAATGTTGAT  
 1381 TTTCAAAGTA AACTTTTACG TAAATATGAT TTGGAAGAAA TAAAGCTCCC  
 CTCAACTATT  
 1441 CGGAAAATAG GTGCTTTTGC TTTTCAATCT AATAACTTGA AATCCTTTGA  
 AGCAAGTGAA  
 1501 GATTTAGAAG AGATTAAAGA GGGAGCCTTT ATGAATAATC GTATTGGAAC  
 TCTAGACTTG  
 1561 AAAGACAAAC TTATCAAAAT AGGTGATGCT GCTTTCCATA TTAATCATAT  
 TTATGCCATT  
 1621 GTTCTTCCAG AATCTGTACA AGAAATAGGA CGTTCAGCTT TTCGACAAAA  
 TGGTGCGCTT

Figure 1 (SEQ ID NO: 1) (continued)

1681 CACCTTATGT TTATCGGAAA TAAGGTTAAA ACAATTGGTG AAATGGCTTT  
TTTATCCAAT  
1741 AAACCTGGAAA GTGTAAATCT CTCTGAGCAA AAACAATTAA AGACAATTGA  
GGTCCAAGCT  
1801 TTTTCGGATA ATGCCCTTAG TGAAGTAGTC TTACCGCCAA ATTTACAGAC  
TATTCGTGAA  
1861 GAGGCTTTCA AAAGGAATCA TTTGAAAGAA GTGAAGGGTT CATCTACATT  
ATCTCAGATT  
1921 ACTTTTAATG CTTTTGATCA AAATGATGGG GACAAACGCT TTGGTAAGAA  
AGTGGTTGTT  
1981 AGGACACATA ATAATTCTCA TATGTTAGCA GATGGTGAGC GTTTTATCAT  
TGATCCAGAT  
2041 AAGCTATCTT CTACAATGGT AGACCTTGAA AAGGTTTTAA AAATAATCGA  
AGGTTTAGAT  
2101 TACTCTACAT TACGTCAGAC TACTCAAACCT CAGTTTAGAG AAATGACTAC  
TGCAGGTAAA  
2161 GCGTTGTTAT CAAAATCTAA CCTCCGACAA GGAGAAAAAC AAAAATTCCT  
TCAAGAAGCA  
2221 CAATTTTTCCT TTGGTCGCGT TGATTTGGAT AAAGCCATAG CTAAAGCTGA  
GAAGGCTTTA  
2281 GTGACCAAGA AGGCAACAAA GAATGGTCAT TTGCTTGAGA GGAGTATTAA  
CAAAGCGGTA  
2341 TTAGCTTATA ATAATAGTGC TATTAAAAAA GCTAATGTGA AGCGCTTGGA  
AAAAGAGTTA  
2401 GACTTGCTGA CAGATTTAGT CGAGGGAAAA GGACCATTAG CGCAAGCTAC  
AATGGTACAA  
2461 GGAGTTTATT TATTAAAGAC GCCTTTACCA TTGCCAGAAT ATTATATCGG  
ATTGAACGTT  
2521 TATTTTGACA AGTCTGGAAA ATTGATTTAT GCACTTGATA TGAGTGATAC  
TATTGGCGAG  
2581 GGACAAAAAG ATGCATATGG TAATCCTATA TTAAATGTTG ACGAGGATAA  
TGAAGGTTAT  
2641 CATACCTTGG CAGTTGCCAC TTTAGCTGAT TATGAAGGTC TTTATATTAA  
AGATATTTTA  
2701 AATAGTTCCC TTGATAAGAT TAAAGCAATA CGCCAGATTC CTTTGGCAAA  
ATATCATAGA  
2761 TTAGGAATTT TCCAAGCTAT CCGAAATGCA GCGGCAGAAG CAGACCGATT  
GCTTCCTAAG  
2821 ACACCTAAGG GGTACCTAAA TGAAGTCCCA AATTATCGTA AAAAACAAAT  
GGAGAAAAAT  
2881 TTAAAACCAG TTGATTATAA AACGCCGATT TTTAATAAGG CTTTACCTAA  
TGAAAAGGTA  
2941 GACGGTGATA GAGCGGCTAA AGGTCATAAT ATAAATGCGG AGACTAATAA  
TTCTGTAGCT  
3001 GTAACACCAA TAAGGTCCGA GCAGCAATTA CATAAGTCAC AGTCTGATGT  
AAATTTACCT  
3061 CAAACAAGTT CTAAAAATAA TTTTATATAC GAGATTCTAG GATACGTTAG  
TTTATGTTTG  
3121 CTTTTCTAG TAACTGCTGG GAAAAAGGA AAACGAGCAA GAAAATAA  
//

Figure 2 (SEQ ID NO: 2)

1 MTKKHLKTLA LALTTVSVVT YSQEVYGLER EESVKQEQTQ SASEDDWFEEDNERKTNVSK  
61 ENSTVDETVS DLFS DGNSNN SSSKTESVVS DPKQVPAKAP EVTQEASNSSNDASKVEVPK  
121 QDTASKKETL ETSTWEAKDF VTRGDTLVGF SKSGINKLSQ TSHLVLP SHAADGTQLTQVA  
181 SFAFTPDKKT AIAEYTSRLG ENGKPSRLDI DQKEIIDEGE IFNAYQLTKLTIPNGYKSIG  
241 QDAFVDNKNI AEVNLPESE TISDYAFAHM SLKQVKLPDN LKVI GELAFFDNQIGGKLYL  
301 PRHLIKLAER AFKSNRIQTV EFLGSKLKVI GEASFQDNNL RNVMLPDGLEKIESEAF TGN  
361 PGDEHYNNQV VLRTRTGQNP HQLATENTYV NPDKSLWRAT PDMDYTKWLEEDFTYQKNSV  
421 TGFSNKGLQK VRRNKNLEIP KQHNGITITE IGDNAFRNVD FQSKTLRKYDLEEIKLPSTI  
481 RKIGAFAFQS NNLKSFEASE DLEEIKEGAF MNNRIGTLDL KDKLIKIGDAAFHINHIIYAI  
541 VLPESVQEIG RSAFRQNGAL HLMFIGNKVK TIGEMAFLSN KLESVNLSEQKQLKTIEVQA  
601 FSDNALSEVV LPPNLQTIRE EAFKR NHLKE VKGSSTLSQI TFNAFDQNDGDKRFGKKVVV  
661 RTHNNSHMLA DGERFIIDPD KLSSTMVDLE KVLKIIIEGLD YSTLRQTTQTQFREMTTAGK  
721 ALLSKSNLRQ GEKQKFLQEA QFFLGRVDLD KAIKA EKAL VTKKATKNGHLLERSINKAV  
781 LAYNNSAIKK ANVKRLEKEL DLLTDLVEGK GPLAQATMVQ GVYLLKTPLPLPEYYIGLNV  
841 YFDKSGKLIY ALDMSDTIGE GQKDAYGNPI LNVDEDNEGY HTLAVATLADYEGLYIKDIL  
901 NSSLDKIKAI RQIPLAKYHR LGIFQAIRNA AAEADRLLPK TPKGYLNEVPNYRKKQMEKN  
961 LKPVDYKTPI FNKALPNEKV DGDRAAKGHN INAETNNSVA VTPIRSEQQLHKSQSDVNLP  
1021 QTSSKNFIY EILGYVSLCL LFLVTAGKKG KRARK\*  
//

## SEQUENCE LISTING

&lt;110&gt; Shire Biochem Inc.

&lt;120&gt; Antigens of Group B Streptococcus and Corresponding DNA Fragments

&lt;130&gt; 74872-81

&lt;150&gt; US 60/287,712

&lt;151&gt; 2001-05-02

&lt;160&gt; 7

&lt;170&gt; PatentIn version 3.0

&lt;210&gt; 1

&lt;211&gt; 3168

&lt;212&gt; DNA

&lt;213&gt; Streptococcus

&lt;400&gt; 1

atgacaaaaa	aacatcttaa	aacgcttgcc	ttggcactta	ctacagtatc	agtagtgaca	60
tacagccagg	aggtatatgg	attagaaaaga	gaggaatcgg	tcaaaacaaga	acaaaccag	120
tcagcttcag	aagatgattg	gttcgaagaa	gataatgaga	ggaaaacaaa	tgtttctaaa	180
gagaattcta	ctgttgatga	aacagttagt	gatttatatt	ctgatggaaa	tagtaataac	240
tctagttcta	aaaccgagtc	agtggtaagt	gaccctaaac	aagtcccaa	agcaaaacca	300
gaggttacac	aagaagcaag	caattctagt	aatgatgcta	gcaaagtaga	agtacaaaa	360
caggatacag	cttcaaaaaa	ggaaactcta	gaaacatcaa	cttgggaggc	aaaagatttc	420
gtaactagag	gggatacttt	agtaggtttt	tcaaaatctg	gaattaataa	gttatctcaa	480
acatcacact	tggttttacc	aagtcatgca	gcagatggaa	ctcaattgac	acaagtagct	540
agctttgctt	ttactccaga	taaaaagacg	gccattgcag	aatatacaag	taggctagga	600
gaaaaatggga	aaccgagtcg	tttagatatt	gatcagaagg	aaattattga	tgagggagaa	660
atattttaatg	cttaccagtt	gactaagctt	actattccaa	atgggtataa	gtctattggg	720
caagatgctt	ttgtggacaa	taagaatatt	gctgagggtta	accttcctga	gagctcgcag	780
actatttcag	actatgcttt	tgtcacatg	tctttaaaac	aagtaaagtt	accagataac	840
ctaaagggtca	ttggagaatt	agcttttttt	gataatcaga	ttggtggtaa	gctttacttg	900
ccacgtcact	tgataaaatt	agcagaacgc	gctttcaaat	ctaactgtat	tcaaacagtt	960
gaatttttgg	gaagtaagct	taagggtata	ggagaagcaa	gttttcaaga	taataatctg	1020
aggaatgtta	tgtctccgga	tggacttgaa	aaaatagaat	cagaagcttt	tacaggaaat	1080
ccaggagatg	aacattacaa	caatcagggt	gtattgcgca	caaggacagg	ccaaaatcca	1140
catcaacttg	cgactgagaa	tacttacgtc	aatccggaca	aatcattgtg	gcgtgcaaca	1200
cctgatatgg	attataccaa	atgggttagag	gaagatttta	cctatcaaaa	aaatagtgtt	1260
acagggtttt	caaataaagg	cttacaaaag	gtaagacgta	ataaaaaact	agaaattcca	1320
aaacaacaca	atgggtattac	tattactgaa	attgggtgata	acgcttttcg	caatggtgat	1380
tttcaaagta	aaactttacg	taaatatgat	ttgggaagaaa	taaagctccc	ctcaactatt	1440
cggaaaatag	gtgcttttgc	ttttcaatct	aataacttga	aatcctttga	agcaagtga	1500
gatttagaag	agattaaaaga	gggagccttt	atgaataatc	gtattggaac	tctagacttg	1560
aaagacaaac	ttatcaaaat	aggtgatgct	gctttccata	ttaatcatat	ttatgccatt	1620
gttcttccag	aatctgtaca	agaaatagga	cgttcagctt	ttcgacaaaa	tggtgcgctt	1680
caccttatgt	ttatcgga	taagggttaa	acaattgggtg	aaatggcttt	tttatccaat	1740
aaactggaaa	gtgtaaatct	ctctgagcaa	aaacaattaa	agacaattga	ggtccaagct	1800
ttttcgata	atgcccttag	tgaagtagtc	ttaccgccaa	atttacagac	tattcgtgaa	1860
gaggctttca	aaaggaatca	tttgaaagaa	gtgaagggtt	catctacatt	atctcagatt	1920
acttttaatg	cttttgatca	aaatgatggg	gacaaacgct	ttggtaagaa	agtggttggt	1980
aggacacata	ataattctca	tatggttagca	gatgggtgagc	gttttatcat	tgatccagat	2040
aagctatctt	ctacaatggg	agaccttgaa	aagggtttta	aaataatcga	aggttttagat	2100
tactctacat	tacgtcagac	tactcaaaact	cagtttagag	aaatgactac	tgacggtaaa	2160
gcgttggtat	caaaatctaa	cctccgacaa	ggagaaaaaac	aaaaattcct	tcaagaagca	2220
caattttttc	ttggtcgcgt	tgattttggat	aaagccatag	ctaaagctga	gaaggcttta	2280
gtgaccaaga	aggcaacaaa	gaatgggtcat	ttgcttgaga	ggagtattaa	caaagcggtta	2340
ttagcttata	ataatagtcg	tattaaaaaa	gctaattgta	agcgcttgga	aaaagagtta	2400
gacttgctga	cagatttagt	cgagggaaaa	ggaccattag	cgcaagctac	aatggtacaa	2460

```

ggagttttatt tattaaagac gcctttacca ttgccagaat attatatcgg attgaacggt 2520
tatttttgaca agtctggaaa attgatttat gcacttgata tgagtgatac tattggcgag 2580
ggacaaaaag atgcatatgg taatcctata ttaaagtgtg acgaggataa tgaaggttat 2640
cataccttgg cagttggcac tttagctgat tatgaaggtc tttatattaa agatatttta 2700
aatagttccc ttgataagat taaagcaata cgccagattc ctttggcaaa atatcataga 2760
ttaggaattt tccaagctat ccgaaatgca gcggcagaag cagaccgatt gcttcctaag 2820
acacctaaagg ggtacctaaa tgaagtccca aattatcgta aaaaacaaat ggagaaaaat 2880
ttaaaccag ttgattataa aacgccgatt ttaataaagg ctttacctaa tgaaaaggta 2940
gacggtgata gagcggctaa aggtcataat ataaatgcgg agactaataa ttctgtagct 3000
gtaacaccaa taaggtcga gcagcaatta cataagtcac agtctgatgt aaatttacct 3060
caaacaagtt ctaaaaataa ttttatatac gagattctag gatacgttag tttatgtttg 3120
cttttctag taactgctgg gaaaaaagga aaacgagcaa gaaaataa 3168

```

```

<210> 2
<211> 1055
<212> PRT
<213> Streptococcus

```

```

<400> 2
Met Thr Lys Lys His Leu Lys Thr Leu Ala Leu Ala Leu Thr Thr Val
1 5 10 15

Ser Val Val Thr Tyr Ser Gln Glu Val Tyr Gly Leu Glu Arg Glu Glu
20 25 30

Ser Val Lys Gln Glu Gln Thr Gln Ser Ala Ser Glu Asp Asp Trp Phe
35 40 45

Glu Glu Asp Asn Glu Arg Lys Thr Asn Val Ser Lys Glu Asn Ser Thr
50 55 60

Val Asp Glu Thr Val Ser Asp Leu Phe Ser Asp Gly Asn Ser Asn Asn
65 70 75 80

Ser Ser Ser Lys Thr Glu Ser Val Val Ser Asp Pro Lys Gln Val Pro
85 90 95

Lys Ala Lys Pro Glu Val Thr Gln Glu Ala Ser Asn Ser Ser Asn Asp
100 105 110

Ala Ser Lys Val Glu Val Pro Lys Gln Asp Thr Ala Ser Lys Lys Glu
115 120 125

Thr Leu Glu Thr Ser Thr Trp Glu Ala Lys Asp Phe Val Thr Arg Gly
130 135 140

Asp Thr Leu Val Gly Phe Ser Lys Ser Gly Ile Asn Lys Leu Ser Gln
145 150 155 160

Thr Ser His Leu Val Leu Pro Ser His Ala Ala Asp Gly Thr Gln Leu
165 170 175

Thr Gln Val Ala Ser Phe Ala Phe Thr Pro Asp Lys Lys Thr Ala Ile
180 185 190

Ala Glu Tyr Thr Ser Arg Leu Gly Glu Asn Gly Lys Pro Ser Arg Leu
195 200 205

Asp Ile Asp Gln Lys Glu Ile Ile Asp Glu Gly Glu Ile Phe Asn Ala
210 215 220

Tyr Gln Leu Thr Lys Leu Thr Ile Pro Asn Gly Tyr Lys Ser Ile Gly
225 230 235 240

```



Gln Asp Ala Phe Val Asp Asn Lys Asn Ile Ala Glu Val Asn Leu Pro  
 245 250 255  
 Glu Ser Leu Glu Thr Ile Ser Asp Tyr Ala Phe Ala His Met Ser Leu  
 260 265 270  
 Lys Gln Val Lys Leu Pro Asp Asn Leu Lys Val Ile Gly Glu Leu Ala  
 275 280 285  
 Phe Phe Asp Asn Gln Ile Gly Gly Lys Leu Tyr Leu Pro Arg His Leu  
 290 295 300  
 Ile Lys Leu Ala Glu Arg Ala Phe Lys Ser Asn Arg Ile Gln Thr Val  
 305 310 315 320  
 Glu Phe Leu Gly Ser Lys Leu Lys Val Ile Gly Glu Ala Ser Phe Gln  
 325 330 335  
 Asp Asn Asn Leu Arg Asn Val Met Leu Pro Asp Gly Leu Glu Lys Ile  
 340 345 350  
 Glu Ser Glu Ala Phe Thr Gly Asn Pro Gly Asp Glu His Tyr Asn Asn  
 355 360 365  
 Gln Val Val Leu Arg Thr Arg Thr Gly Gln Asn Pro His Gln Leu Ala  
 370 375 380  
 Thr Glu Asn Thr Tyr Val Asn Pro Asp Lys Ser Leu Trp Arg Ala Thr  
 385 390 395 400  
 Pro Asp Met Asp Tyr Thr Lys Trp Leu Glu Glu Asp Phe Thr Tyr Gln  
 405 410 415  
 Lys Asn Ser Val Thr Gly Phe Ser Asn Lys Gly Leu Gln Lys Val Arg  
 420 425 430  
 Arg Asn Lys Asn Leu Glu Ile Pro Lys Gln His Asn Gly Ile Thr Ile  
 435 440 445  
 Thr Glu Ile Gly Asp Asn Ala Phe Arg Asn Val Asp Phe Gln Ser Lys  
 450 455 460  
 Thr Leu Arg Lys Tyr Asp Leu Glu Glu Ile Lys Leu Pro Ser Thr Ile  
 465 470 475 480  
 Arg Lys Ile Gly Ala Phe Ala Phe Gln Ser Asn Asn Leu Lys Ser Phe  
 485 490 495  
 Glu Ala Ser Glu Asp Leu Glu Glu Ile Lys Glu Gly Ala Phe Met Asn  
 500 505 510  
 Asn Arg Ile Gly Thr Leu Asp Leu Lys Asp Lys Leu Ile Lys Ile Gly  
 515 520 525  
 Asp Ala Ala Phe His Ile Asn His Ile Tyr Ala Ile Val Leu Pro Glu  
 530 535 540  
 Ser Val Gln Glu Ile Gly Arg Ser Ala Phe Arg Gln Asn Gly Ala Leu  
 545 550 555 560  
 His Leu Met Phe Ile Gly Asn Lys Val Lys Thr Ile Gly Glu Met Ala  
 565 570 575

Phe Leu Ser Asn Lys Leu Glu Ser Val Asn Leu Ser Glu Gln Lys Gln  
 580 585 590  
 Leu Lys Thr Ile Glu Val Gln Ala Phe Ser Asp Asn Ala Leu Ser Glu  
 595 600 605  
 Val Val Leu Pro Pro Asn Leu Gln Thr Ile Arg Glu Glu Ala Phe Lys  
 610 615 620  
 Arg Asn His Leu Lys Glu Val Lys Gly Ser Ser Thr Leu Ser Gln Ile  
 625 630 635 640  
 Thr Phe Asn Ala Phe Asp Gln Asn Asp Gly Asp Lys Arg Phe Gly Lys  
 645 650 655  
 Lys Val Val Val Arg Thr His Asn Asn Ser His Met Leu Ala Asp Gly  
 660 665 670  
 Glu Arg Phe Ile Ile Asp Pro Asp Lys Leu Ser Ser Thr Met Val Asp  
 675 680 685  
 Leu Glu Lys Val Leu Lys Ile Ile Glu Gly Leu Asp Tyr Ser Thr Leu  
 690 695 700  
 Arg Gln Thr Thr Gln Thr Gln Phe Arg Glu Met Thr Thr Ala Gly Lys  
 705 710 715 720  
 Ala Leu Leu Ser Lys Ser Asn Leu Arg Gln Gly Glu Lys Gln Lys Phe  
 725 730 735  
 Leu Gln Glu Ala Gln Phe Phe Leu Gly Arg Val Asp Leu Asp Lys Ala  
 740 745 750  
 Ile Ala Lys Ala Glu Lys Ala Leu Val Thr Lys Lys Ala Thr Lys Asn  
 755 760 765  
 Gly His Leu Leu Glu Arg Ser Ile Asn Lys Ala Val Leu Ala Tyr Asn  
 770 775 780  
 Asn Ser Ala Ile Lys Lys Ala Asn Val Lys Arg Leu Glu Lys Glu Leu  
 785 790 795 800  
 Asp Leu Leu Thr Asp Leu Val Glu Gly Lys Gly Pro Leu Ala Gln Ala  
 805 810 815  
 Thr Met Val Gln Gly Val Tyr Leu Leu Lys Thr Pro Leu Pro Leu Pro  
 820 825 830  
 Glu Tyr Tyr Ile Gly Leu Asn Val Tyr Phe Asp Lys Ser Gly Lys Leu  
 835 840 845  
 Ile Tyr Ala Leu Asp Met Ser Asp Thr Ile Gly Glu Gly Gln Lys Asp  
 850 855 860  
 Ala Tyr Gly Asn Pro Ile Leu Asn Val Asp Glu Asp Asn Glu Gly Tyr  
 865 870 875 880  
 His Thr Leu Ala Val Ala Thr Leu Ala Asp Tyr Glu Gly Leu Tyr Ile  
 885 890 895  
 Lys Asp Ile Leu Asn Ser Ser Leu Asp Lys Ile Lys Ala Ile Arg Gln  
 900 905 910

Ile Pro Leu Ala Lys Tyr His Arg Leu Gly Ile Phe Gln Ala Ile Arg  
 915 920 925

Asn Ala Ala Ala Glu Ala Asp Arg Leu Leu Pro Lys Thr Pro Lys Gly  
 930 935 940

Tyr Leu Asn Glu Val Pro Asn Tyr Arg Lys Lys Gln Met Glu Lys Asn  
 945 950 955 960

Leu Lys Pro Val Asp Tyr Lys Thr Pro Ile Phe Asn Lys Ala Leu Pro  
 965 970 975

Asn Glu Lys Val Asp Gly Asp Arg Ala Ala Lys Gly His Asn Ile Asn  
 980 985 990

Ala Glu Thr Asn Asn Ser Val Ala Val Thr Pro Ile Arg Ser Glu Gln  
 995 1000 1005

Gln Leu His Lys Ser Gln Ser Asp Val Asn Leu Pro Gln Thr Ser  
 1010 1015 1020

Ser Lys Asn Asn Phe Ile Tyr Glu Ile Leu Gly Tyr Val Ser Leu  
 1025 1030 1035

Cys Leu Leu Phe Leu Val Thr Ala Gly Lys Lys Gly Lys Arg Ala  
 1040 1045 1050

Arg Lys  
 1055

<210> 3  
 <211> 34  
 <212> DNA  
 <213> Artificial

<220>  
 <223> primer

<400> 3  
 gcgcccattgg tgcaggaggt atatgggttag aaag

34

<210> 4  
 <211> 30  
 <212> DNA  
 <213> Artificial

<220>  
 <223> primer

<400> 4  
 ggtggatccg agaaaggctt tattgtaatg

30

<210> 5  
 <211> 36  
 <212> DNA  
 <213> Artificial

<220>  
 <223> primer

<400> 5  
catattaatt gcggccgctt ttcttgctcg ttttcc

36

<210> 6  
<211> 34  
<212> DNA  
<213> Artificial

<220>  
<223> primer

<400> 6  
cgttggatcc tcaggaggta tatggattag aaag

34

<210> 7  
<211> 31  
<212> DNA  
<213> Artificial

<220>  
<223> primer

<400> 7  
catcgtcgac ttatttttctt gctcgttttc c

31